

LANCS-TH/9821
 hep-ph/9810320
 (October 1998)

Inflation with TeV-scale gravity needs supersymmetry

David H. Lyth

*Department of Physics,
 Lancaster University,
 Lancaster LA1 4YB. U. K.*

E-mail: d.lyth@lancaster.ac.uk

Abstract

Allowing for the possibility of large extra dimensions, the fundamental Planck scale M could be anywhere in the range $\text{TeV} \lesssim M \lesssim M_{\text{P}}$, where $M_{\text{P}} = 2.4 \times 10^{18} \text{ GeV}$ is the four-dimensional Planck scale. If $M \sim \text{TeV}$, quantum corrections would not destabilize the Higgs mass even if there were no supersymmetry. But we point out that supersymmetry must in fact be present, if there is an era of cosmological inflation, since during such an era the inflaton mass satisfies $m \ll M^2/M_{\text{P}} = 10^{-15}(M/\text{TeV})$ and supersymmetry will be needed to protect it. If the inflation hypothesis is accepted, there is no reason to think that Nature has chosen the low value $M \sim \text{TeV}$, however convenient that choice might have been for the next generation of collider experiments.

1. There is a large hierarchy, $m_{\text{H}}/M_{\text{P}} \sim 10^{-15}$, between the mass $m_{\text{H}} \sim 1 \text{ TeV}$ of the Standard Model Higgs field and the Planck scale $M_{\text{P}} \equiv (8\pi G)^{-1/2} \simeq 10^{18} \text{ GeV}$. This makes it difficult to understand the existence of the Higgs field, because in a generic field theory valid up to the Planck scale every elementary scalar field will have a mass of order M_{P} . To be precise, the mass will be given by $m^2 = m_0^2 + \sum \Delta_i + \dots$, where the first term is the tree-level value and the Δ_i are one-loop contributions of order $(\lambda_i M_{\text{P}})^2$, with λ_i the strength of the interaction. To avoid this disaster, it is usual to invoke supersymmetry which automatically cancels the one-loop contributions to sufficient accuracy.¹

An alternative proposal [2] is to place the fundamental Planck scale M in the TeV region. This makes the one-loop contributions of order $(\lambda_i M)^2$ and avoids fine-tuning.

¹The alternatives are an accidental cancellation, perhaps to be understood anthropically [1], or a composite Higgs such as occurs in technicolor theories. It is difficult to construct composite theories that are consistent with observation.

The low value of M is achieved by invoking extra space dimensions with large compactification radius. If there are n extra space dimensions with compactification radius R , and M is the fundamental Planck scale in $4 + n$ spacetime dimensions, the Planck scale that we observe is given by

$$M_{\text{P}}^2 \sim R^n M^{2+n}. \quad (1)$$

Einstein or Newtonian gravity holds on scales $\gtrsim R$, but is modified on smaller scales. Putting $M \sim 1 \text{ TeV}$ and $n \geq 2$ gives $R \gtrsim 1 \text{ mm}$, which is allowed by observation since the law of gravity is unknown on scales $\lesssim 1 \text{ cm}$. One can envisage more complicated schemes, where the compactified dimensions have different radii, but in all cases the biggest dimension must be $R \lesssim 1 \text{ cm}$.

In this scheme, the hierarchy $m_{\text{H}}/M_{\text{P}}$ is replaced by the hierarchy R^{-1}/M , which is of a different type and might be easier to understand. Investigations reported so far [3] seem to suggest that the scheme is viable, with no need for supersymmetry in the field theory that contains the Standard Model.² As I now explain, this apparent success disappears as soon as one tries to construct a model of inflation, that is presumably necessary to generate structure in the Universe. A more detailed investigation will be reported elsewhere [5].

2. We are concerned with the cosmology of the observable Universe. The Universe is modeled as a practically homogeneous and isotropic fluid, with the distance between comoving fluid elements proportional to a universal scale factor $a(t)$. The evolution of the scale factor is given by the Friedmann equation, which assuming spatial flatness is³

$$3H^2 = \rho/M_{\text{P}}^2, \quad (2)$$

and the continuity equation

$$\dot{\rho} = -3H(\rho + P). \quad (3)$$

Here $H = \dot{a}/a$ is the Hubble parameter, ρ is the energy density and P is the pressure.

The Friedmann and continuity equations are consequences of Einstein's field equation, and are valid provided that all relevant quantities are smoothed on a comoving distance scale $\gg R$. The cosmic fluid, which is the subject of cosmology, is defined at a given epoch only after such smoothing. (There is no question of 'modifying the Friedmann equation on short distance scales' since we are dealing with a universal scale factor.)

The history of the Universe begins at some energy density $\lesssim M^4$, where $M \sim \text{TeV}$ is the fundamental Planck scale. In particular, the potential $V \simeq \rho$ during inflation satisfies

$$V(\phi) \lesssim M^4. \quad (4)$$

²Other works, for example [4], have explored the possibility of large extra dimensions within the context of supersymmetry.

³For simplicity, I am assuming that M_{P} during inflation has its present value. With M declared fixed (the choice of energy unit) this amounts to saying that R has its present value. The more general possibility is considered elsewhere [5].

The vacuum fluctuation of the inflaton field ϕ , that is supposed to be the origin of large scale structure, is generated on each comoving scale a/k at the epoch of horizon exit $k = aH$. We therefore require the Hubble distance H^{-1} to be much bigger than the radius of the internal dimensions, $HR \ll 1$. Because of Eq. (4), this is not a very severe restriction. Since M^{-1} is the smallest distance that makes sense in the context of quantum gravity, we must have $RM \gtrsim 1$. Then Eqs. (4) and (2) give $H^{-1} \gtrsim 1 \text{ mm}$, whereas observation requires $R \lesssim 1 \text{ cm}$. Irrespective of observation, Eqs. (1) and (4) require $(RH)^2 \lesssim 1$ if $n \geq 2$ [6]. (Indeed, they give $(RH)^2 \sim (MR)^{2-n}$.) The inclusion of additional, smaller dimensions only strengthens this result.

If m is the inflaton mass during inflation,⁴ the potential is of the form

$$V = V_0 \pm \frac{1}{2}m^2\phi^2 + \dots \quad (5)$$

The dots represent additional terms, which might come from a variety of sources [6] (higher powers of ϕ representing interaction terms in the tree-level potential, logarithmic terms representing loop corrections etc.). In order to generate the nearly scale-invariant primordial curvature perturbation, that is presumed to be the origin of large scale structure, one should have [6]

$$\left| \frac{M_{\text{P}}^2 V''}{V} \right| \ll 1, \quad (6)$$

while cosmological scales are leaving the horizon.

Since each term has a different ϕ dependence, and ϕ usually varies significantly over cosmological scales, there is hardly likely to be an accurate cancellation between terms. Then, discounting the possibility $\phi \gg M_{\text{P}}$ which would surely place the field theory out of control, the flatness condition has to be satisfied by each term individually, with V dominated by the constant term V_0 . In particular, the mass has to satisfy $m^2 \ll V_0/M_{\text{P}}^2$. Remembering that $V_0 \lesssim M^4$ this becomes [8] $m^2 \ll M^4/M_{\text{P}}^2 \sim (1 \text{ mm})^{-2}$, or

$$\frac{m}{M} \ll \frac{M}{M_{\text{P}}} \sim 10^{-15}. \quad (7)$$

To summarise, taking the fundamental Planck scale M to be of order 1 TeV removes the hierarchy between the Higgs mass and M , at the expense of introducing at least the *same* hierarchy between the inflaton mass and M . To protect the inflaton mass from quantum corrections, supersymmetry is needed, just as it was needed to protect the Higgs mass in the case $M = M_{\text{P}}$.

⁴During inflation, the ‘vacuum’ for quantum field theory is defined by the values of the inflaton and other relevant fields, which may be different from their true vacuum values. In hybrid inflation models [7] a field ψ has a coupling like $\psi^2\phi^2$, which holds it at the origin during inflation. Afterwards, ψ acquires its true vacuum value, which can give the inflaton a large mass in the vacuum. As a result, the quantum field theory containing Standard Model particles and the inflaton could have all scalar masses of order TeV *in the true vacuum*. But we need a sensible quantum field theory also during inflation.

In order to reheat the Universe, the inflaton must have significant couplings (not necessarily tree-level) with Standard Model particles. As a result, these particles should belong to the same supersymmetric field theory as the inflaton.⁵

If one accepts the hypothesis of cosmological inflation, the original motivation [2] for considering $M \sim 1 \text{ TeV}$ is now removed, and there seems to be no reason why Nature should have chosen this value. Still, one may choose to explore that possibility, either because it will be accessible to observation in the foreseeable future or because a lot of effort has been invested in it.

3. In that case, one might ask whether a viable model of inflation can be constructed. It is easy enough to write down a potential $V(\phi) \lesssim M^4$, valid during slow-roll inflation, that gives the correct curvature perturbation $\delta_H = 1.9 \times 10^{-5}$ on COBE scales and a spectral index within the observed band $|n - 1| < 0.2$. Take, for instance, the potential

$$V = V_0 - \frac{1}{4}\lambda\phi^4 + \dots, \quad (8)$$

with the additional terms negligible during slow-roll inflation. Let us assume that

$$\phi_{\text{end}}/\phi_{\text{COBE}} \gg 1, \quad (9)$$

where ϕ_{end} is the end of slow-roll inflation and ϕ_{COBE} is the epoch when COBE scales leave the horizon. Using well-known formulas [6], the COBE constraint is $\lambda \sim 10^{-14}(30/N)^2$ independently of V_0 , and $n - 1 = -3/N$. Here N is the number of e -folds after COBE scales leave the horizon, given (discounting thermal inflation and late-decaying particles) by

$$N \simeq 30 - \ln(1 \text{ TeV}/V^{1/4}) - \frac{1}{3} \ln(V^{1/4}/T_{\text{reh}}). \quad (10)$$

Moreover, $\phi_{\text{COBE}} \simeq 10^{-5} \text{ GeV}$, so the initial assumption Eq. (9) is not very restrictive.

One might also wish to impose the constraint $\phi \lesssim M \sim \text{TeV}$; for instance this might be necessary to have control over non-renormalizable terms if they are of order ϕ^d/M^{d-4} , or to have control over the running of couplings and masses in a renormalizable theory. In the above example this constraint no problem, but in general it is a severe restriction as will be discussed elsewhere [5].

4. It is generally accepted that a viable cosmology should begin with an era of inflation, to set suitable initial conditions for the subsequent hot big bang and in particular to provide an origin for structure. We have argued that the inflaton mass during inflation has to satisfy $m \ll (M^2/M_{\text{P}}) = 10^{-15}(M/\text{TeV})$, and that supersymmetry should be invoked to stabilize this mass (or the masses of scalar fields produced after inflation). A more detailed investigation [5] supports this conclusion. If one accepts it, along with the need for inflation, one concludes that there is no reason to think that Nature has chosen $M \sim \text{TeV}$, however convenient such a choice might have been for the next generation of collider experiments.

⁵By contrast, the Kaluza-Klein tower of scalar particles, associated with the extra dimensions, need not be considered as part of the same theory since they will have very weak coupling. For the same reason, their masses are presumably not destabilized by loop corrections.

Acknowledgements

This work was initiated at CERN. I am grateful to CERN for support, and to Steve Abel, Karim Benakli, Keith Dienes, Tony Gerghetta, John March-Russell and Toni Riotto for useful discussions and correspondence.

References

- [1] V. Agrawal, S. M. Barr, J. F. Donoghue and D. Seckel, *Phys. Rev.* **D57** (1998) 5480.
- [2] N. Arkani-Hamed, S. Dimopoulos and G. Dvali, hep-ph/9803315, to appear in *Phys. Lett.* **B**.
- [3] I. Antoniadis, N. Arkani-Hamed, S. Dimopoulos and G. Dvali, hep-ph/9804398, to appear in *Phys. Lett.* **B**; N. Arkani-Hamed, S. Dimopoulos and G. Dvali, hep-ph/9807344; N. Arkani-Hamed, S. Dimopoulos and J. March-Russell, hep-th/9809124; P. C. Argyres, S. Dimopoulos and J. March-Russell, hep-th/9808138; R. Sundrum, hep-ph/9807348.
- [4] K.R. Dienes, E. Dudas and T. Gherghetta, hep-ph/9803466 and hep-ph/9806292; K.R. Dienes, E. Dudas, T. Gherghetta and A. Riotto, hep-ph/9809406. G. Shiu and S. H. H. Tye, hep-th/9805157. R. Sundrum, hep-ph/9805471.
- [5] D. H. Lyth, in preparation.
- [6] D. H. Lyth and A. Riotto, hep-ph/9807278, to appear in Phys. Reports.
- [7] A. D. Linde, *Phys. Lett.* **B259** (1991) 38.
- [8] K. Benakli and S. Davidson, hep-ph/9810280.
- [9] P. Horava and E. Witten, *Nucl. Phys.* **B475** (1996) 94; *Nucl. Phys.* **B460** (1996) 506.
- [10] J.D. Lykken, *Phys. Rev.* **D54** (1996) 3693.